

1.

Morak M, Schaefer K, Steinke-Lange V, et al. Full-length transcript amplification and sequencing as universal method to test mRNA integrity and biallelic expression in mismatch repair genes. *Eur J Hum Genet.* July 2019. doi:[10.1038/s41431-019-0472-8](https://doi.org/10.1038/s41431-019-0472-8)

2.

Hallermayr A, Graf J, Koehler U, et al. Extending the critical regions for mutations in the non-coding gene RNU4ATAC in another patient with Roifman Syndrome. *Clin Case Rep.* 2018;6(11):2224-2228. doi:[10.1002/ccr3.1830](https://doi.org/10.1002/ccr3.1830)

3.

Oliveira J, Gruber A, Cardoso M, et al. LAMA2 gene mutation update: Toward a more comprehensive picture of the laminin-alpha2 variome and its related phenotypes. *Hum Mutat.* 2018;39(10):1314-1337. doi:[10.1002/humu.23599](https://doi.org/10.1002/humu.23599)

4.

Morak M, Ibisler A, Keller G, et al. Comprehensive analysis of the MLH1 promoter region in 480 patients with colorectal cancer and 1150 controls reveals new variants including one with a heritable constitutional MLH1 epimutation. *J Med Genet.* 2018;55(4):240-248. doi:[10.1136/jmedgenet-2017-104744](https://doi.org/10.1136/jmedgenet-2017-104744)

5.

Morak M, Kasbauer S, Kerscher M, et al. Loss of MSH2 and MSH6 due to heterozygous germline defects in MSH3 and MSH6. *Fam Cancer.* 2017;16(4):491-500. doi:[10.1007/s10689-017-9975-z](https://doi.org/10.1007/s10689-017-9975-z)

6.

Adam R, Spier I, Zhao B, et al. Exome Sequencing Identifies Biallelic MSH3 Germline Mutations as a Recessive Subtype of Colorectal Adenomatous Polyposis. *Am J Hum Genet.* 2016;99(2):337-351. doi:[10.1016/j.ajhg.2016.06.015](https://doi.org/10.1016/j.ajhg.2016.06.015)

7.

Spier I, Kerick M, Drichel D, et al. Exome sequencing identifies potential novel candidate genes in patients with unexplained colorectal adenomatous polyposis. *Fam Cancer.* 2016;15(2):281-288. doi:[10.1007/s10689-016-9870-z](https://doi.org/10.1007/s10689-016-9870-z)

8.

Spier I, Drichel D, Kerick M, et al. Low-level APC mutational mosaicism is the underlying cause in a substantial fraction of unexplained colorectal adenomatous polyposis cases. *J Med Genet.* 2016;53(3):172-179. doi:[10.1136/jmedgenet-2015-103468](https://doi.org/10.1136/jmedgenet-2015-103468)

9.

Wilhelm F, Kassner F, Schmid G, et al. Phosphatidylinositol 3-kinase (PI3K) signalling regulates insulin-like-growth factor binding protein-2 (IGFBP-2) production in human adipocytes. *Growth Horm IGF Res.* 2015;25(3):115-120. doi:[10.1016/j.gHIR.2015.03.003](https://doi.org/10.1016/j.gHIR.2015.03.003)

10.

Horpaopan S, Spier I, Zink AM, et al. Genome-wide CNV analysis in 221 unrelated patients and targeted high-throughput sequencing reveal novel causative candidate genes for colorectal adenomatous polyposis. *Int J Cancer*. 2015;136(6):E578-589. doi:[10.1002/ijc.29215](https://doi.org/10.1002/ijc.29215)

11.

Morak M, Heidenreich B, Keller G, et al. Biallelic MUTYH mutations can mimic Lynch syndrome. *Eur J Hum Genet*. 2014;22(11):1334-1337. doi:[10.1038/ejhg.2014.15](https://doi.org/10.1038/ejhg.2014.15)

12.

Schmid GL, Kassner F, Uhlig HH, et al. Sirolimus treatment of severe PTEN hamartoma tumor syndrome: case report and in vitro studies. *Pediatr Res*. 2014;75(4):527-534. doi:[10.1038/pr.2013.246](https://doi.org/10.1038/pr.2013.246)

13.

Rocchi L, Braz C, Cattani S, et al. Escherichia coli-cloned CFTR loci relevant for human artificial chromosome therapy. *Hum Gene Ther*. 2010;21(9):1077-1092. doi:[10.1089/hum.2009.225](https://doi.org/10.1089/hum.2009.225)

14.

Morak M, Laner A, Scholz M, Madorf T, Holinski-Feder E. Report on de-novo mutation in the MSH2 gene as a rare event in hereditary nonpolyposis colorectal cancer. *Eur J Gastroenterol Hepatol*. 2008;20(11):1101-1105. doi:[10.1097/MEG.0b013e328305e185](https://doi.org/10.1097/MEG.0b013e328305e185)

15.

Dames P, Laner A, Maucksch C, Aneja MK, Rudolph C. Targeting of the glucocorticoid hormone receptor with plasmid DNA comprising glucocorticoid response elements improves nonviral gene transfer efficiency in the lungs of mice. *J Gene Med*. 2007;9(9):820-829. doi:[10.1002/jgm.1082](https://doi.org/10.1002/jgm.1082)

16.

Huth S, Hoffmann F, von Gersdorff K, et al. Interaction of polyamine gene vectors with RNA leads to the dissociation of plasmid DNA-carrier complexes. *J Gene Med*. 2006;8(12):1416-1424. doi:[10.1002/jgm.975](https://doi.org/10.1002/jgm.975)

17.

Klink D, Schindelhauer D, Laner A, et al. Gene delivery systems--gene therapy vectors for cystic fibrosis. *J Cyst Fibros*. 2004;3 Suppl 2:203-212. doi:[10.1016/j.jcf.2004.05.042](https://doi.org/10.1016/j.jcf.2004.05.042)